

# STUDIES OF ELECTROCHEMICAL SYSTEMS WITH RUNNING FARADAIC REACTIONS BY IN SITU SPEM

B. Bozzini<sup>1</sup>, C. Mele<sup>1</sup>, E. Tondo<sup>1</sup>, M. Amati<sup>2</sup>, L. Gregoratti<sup>2</sup>, M. Kazemian Abyaneh<sup>2</sup>, M. Kiskinova<sup>2</sup>

<sup>1</sup>*I.N.F.M. Unità di Lecce, Università di Lecce, via per Monteroni, I-73100 Lecce, Italy*

<sup>2</sup>*Sincrotrone Trieste SCpA, SS14-km163.5 in Area Science Park, 34149 Trieste, Italy*

Advanced studies in anodic and cathodic electrochemistry require dynamic position-dependent compositional, morphological and structural information at the submicrometric and mesoscopic scales (0.1÷25  $\mu\text{m}$ ), regarding the reactive interface involved in faradaic processes. The zone-plate based scanning photoemission microscopy, allowing different detection modes and variable fields of view, currently gives the possibility of: (i) applying external potentials, (ii) probing samples with rough surfaces and (iii) dosing gases in the analysis chamber: electrochemical science and technology can greatly profit of these capabilities,.

This presentation will address three key issues in the field of in situ electrochemical SPEM: (1) briefly review the state of the art regarding in situ electrochemical SPEM and XPS experiments carried out with both synchrotron and conventional sources; (2) report original SPEM results obtained by the authors at the ESCAmicroscopy beamline at Elettra and (3) discuss the perspectives of SPEM-based approaches for the in situ investigation of cutting-edge electrochemical materials science problems.




Figure 1




Figure 2

Figure 1: Dynamic Ni 3p XPS spectra during reduction at (A)  $-625 \text{ mV}_{\text{cell}}$  and (B)  $-3000 \text{ mV}_{\text{cell}}$ . Red mark: elemental Ni, light blue mark: Ni(II), blue mark Ni satellite peak.

Figure 2: SPEM images of (A) cathode (Ni electrode on the left,  $30 \times 256 \times 2 \mu\text{m}$ ) and (B) anode (Ni electrode on the right  $64 \times 256 \times 2 \mu\text{m}$ ), at the Ni 3p energy (611 eV), under electrochemical polarisation ( $-625 \text{ mV}_{\text{cell}}$ ).